

# POST TREATMENT INFORMATION BOTULINUM TOXIN A

- Do not lie down, or rub or massage the treated area for 4 hours after treatment
- You can help activate the Botox by gently exercising the muscles that have been treated, i.e. frown and smile as frequently as possible.
- Some patients experience slight discomfort around the treatment area for a short time after treatment.
- Bruising can be avoided or minimized by avoiding aspirin and alcohol for 24 hours pre & post treatment
- You may experience temporary swelling or bruising around the site of treatment. This will usually settle within a day or two. Some patients can experience a headache following treatment for which you may take paracetamol.
- The results are permanent and the duration will differ from patient to patient.
- The treatment normally takes 4-10 days to take effect.
- Side effects if they occur are usually temporary and mild to moderate.

## The most common reported side effects:

- Tenderness
- Bruising
- Temporary Headache
- Drooping of the eyelid or brow
- Excessive watering of the eye
- Mild inflammation of the surface of the eye

## Less common side effects include:

- Inflammation of the surface of the eye
- Eversion or inversion of the eyelids
- Light sensitivity/Double vision
- Dizziness, Rash, Tiredness
- Facial weakness/droop/asymmetry
- Allergic reaction

Should you experience any of these symptoms following treatment please contact your practitioner **Olha Vorodyukhina** on **07921851015**

**WE ALSO OFFER:** Dermal Fillers / Lip Augmentation / Chemical Peels / Dermaroller Treatment / Teeth Whitening. **Please ask your practitioner for details.**